



# International Journal of Advanced Pharmaceutics

[www.ijapjournal.com](http://www.ijapjournal.com)

## NAIL DRUG DELIVERY SYSTEM-A NOVEL APPROACHES FOR DRUG DELIVERY SYSTEM

**G.Poornima<sup>\*</sup>, K.Sandhya Rani, B.Brahmaiah, K.Padmaja, Ch.Pratyusha**

Department of Pharmaceutics, Priyadarshini Institute of Pharmaceutical Education & Research(PIPER), 5<sup>th</sup> Mile, Pulladigunta, Kornepadu (V), Vatticherukuru (M), Guntur-522017, Andhra Pradesh, India.

### ABSTRACT

Topical therapy is highly desirable in treating nail disorders due to its localized effects, which results in minimal adverse systemic events and possibly improved adherence. However, the effectiveness of topical therapies is limited by minimal drug permeability through the nail plate. Infections of foot and hand nails by fungi are a very common condition in millions of people. They account for about half of all nail disorders and are estimated to occur in over 10% of the population. Such infections may be difficult to treat, and currently prescribed oral antifungal medications may cause side effects ranging from skin rashes to liver damage. Other treatment modalities include the use of antifungal lacquers and topical medications. Topical trans-nail delivery of antifungal drugs is limited by several physicochemical and physiological factors. Use of chemical permeation enhancers has been a common approach for enhancing trans-nail delivery of drugs. The potential of physical permeation enhancement techniques has been found to be higher than the potential of chemical permeation enhancers in transdermal delivery of hydrophilic drugs and macromolecular therapeutic agents. However, application of physical permeation enhancement techniques has not been explored for trans-nail drug delivery. In the current work, iontophoresis was applied across human nail in vitro to assess its efficiency in enhancing drug delivery. This new therapy may reduce the need for hazardous systemic administration of oral antifungal drugs for nail infections. A novel ultrasound-mediated drug delivery system has been developed for treatment of a nail fungal disorder (onychomycosis) by improving delivery to the nail bed using ultrasound to increase the permeability of the nail. The slip-in device consists of ultrasound transducers and drug delivery compartments above each toenail. The device is connected to a computer, where a software interface allows users to select their preferred course of treatment.

**Keywords:** Topical Trans-Nail Delivery, Human Nail, Antifungal Drugs, Iontophoresis.

### INTRODUCTION

The nail is horny structure. Nail plate is responsible for penetration of drug across it. As it is hard enough the penetration becomes difficult, only a fraction of topical drug penetrates across it. Hence the effective therapeutic concentration is not achieved. The nail plate may appear abnormal as a result of decreased glow. It's involvement of nail bed, reduction of blood supply, physical or chemical features of nail bed. As a result variety of diseases occurs [1]. These diseases can be cured by achieving desired therapeutic concentration of drug by nail drug delivery system. Human nails do not have and decorative role, but can also be considered as an alternative

pathway for drug. Delivery, especially in nail diseases such as onychomycosis or psoriasis. These nail diseases are widely spread in the population, particularly among elderly and immune compromised patients. Oral therapies are accompanied by systemic side effects and drug interactions, while topical therapies are limited by the low permeation rate through the nail plate. For the successful treatment of nail disease the applied active drug must permeate through the dense keratinized nail plate and reach deeper layers, the nail bed and the nail matrix. Studies conducted on the human skin elucidated its structure, functions, and its permeability for some substances, but very little is known about skin derivate, the nail, and the

---

Corresponding Author:- **G.Poornima** Email:- brahmaiahmph@gmail.com

properties of nail keratin. The purpose of this work is to improve the understanding of physicochemical parameters that influence drug permeation through the nail plate in order to treat not only topical nail diseases but also to consider the possibility to reach systemic circulation and neighboring target sites [2]. The purpose of this review is to explore the difficulties in penetration of drug across nail plate & enhancement of bioavailability of antifungal drug. The evidence suggests that a key to successful treatment of fungal diseases by topical antifungal product lies in ineffectively overcoming the nail barrier. Current topical treatments have limited therapeutic effectiveness possibly because they can not sufficiently penetrate in the nail plate to transport a therapeutically sufficient quantity of antifungal drug to the target sites to eradicate the protection. Also the analysis of the drug's penetration is a difficult task. The topical therapy of nail diseases, especially of onychomycosis, and to a smaller extent, of nail psoriasis, is desirable to avoid the side effects associated with their systemic therapy, to increase patient compliance and reduce the cost of treatment. Systemic therapy is however the mainstay of treatment due to the poor permeability of the nail plate to topically applied drugs. For effective topical therapy, fungal drug permeation must be enhanced [3]. This can be achieved by disrupting the nail plate using physical techniques or chemical agents. Alternatively, drug permeation into the intact nail plate may be encouraged, for example, by iontophoresis or by formulating the drug within a vehicle which enables high drug partition out of the vehicle and into the nail plate [4]. The physical techniques (manual and electrical nail abrasion, acid etching, ablation by lasers, microporation, application of low-frequency ultrasound and electric currents) and chemicals (thiols, sulphites, hydrogen peroxide, urea, water, enzymes) that have shown fungal enhancer activity. The human nail can be afflicted by several disease states including paronychia, psoriasis and infections due to bacteria, viruses or fungi. Whilst rarely life threatening, these generate self-consciousness and psychological stress [4]. Approximately 50% of all problems result from fungal infections, onychomycoses, and the prevalence of these may be as high as 27% in Europe and 10% in the United States. There are many treatment regimens, but the most common involves oral dosing with antifungal agents such as terbinafine or itraconazole. Experimental techniques for investigation of the penetration and distribution of chemicals into and through the nail plate demonstrated that it is possible to deliver drugs to the nail following topical application and led to the development of newer more effective topical products and regimens for treatment of onychomycoses and other nail diseases. A novel ultrasound-mediated drug delivery system has been developed for treatment of a nail fungal disorder (onychomycosis) by improving delivery to the nail bed using ultrasound to increase the permeability of the nail [5].

## ANATOMY OF NAIL

### Parts of the nail

The matrix (sometimes called the matrix unguis, keratogenous membrane, nail matrix, or onychostroma) is the tissue (or germinal matrix) which the nail protects,] the part of the nail bed that rests beneath the nail and contains nerves, lymph and blood vessels [5]. The matrix is responsible for the producing cells that become the nail plate. The width and thickness of the nail plate is determined by the size, length, and thickness of the matrix, while the shape of the fingertip itself shows if the nail plate is flat, arched or hooked. The matrix will continue to grow as long as it receives nutrition and remains in a healthy condition. As new nail plate cells are made, they push older nail plate cells forward and in these ways older cells become compressed, flat, and translucent. This makes the capillaries in the nail bed below visible, resulting in a pink color. The lunula (or simply "the moon") is the visible part of the matrix, the whitish crescent shaped base of the visible nail. The lunula can be seen as the largest in the thumb and often is not present in the little finger. The nail bed is the skin beneath the nail plate. Like all skin, it is made of two types of tissues: the deeper dermis, the living tissue fixed to the bone which includes capillaries and glands, [ and the superficial epidermis, the layer just beneath the nail plate which moves forward with the plate. The epidermis is attached to the dermis by tiny longitudinal "grooves known as matrix crests (cristae matricis unguis). During old age, the plate thins and these grooves are more visible <sup>7</sup>The nail sinus (sinus unguis) is where the nail root is inserted. The nail root (radix unguis) is the part of nail situated in the nail sinus, i.e. the base of the nail under beneath the skin. It originates from the actively growing tissue below, the matrix. The nail plate (corpus unguis) is the actual nail, made of translucent keratin protein. Several layers of dead, compacted cells cause the nail to be strong but flexible. Its (transversal) shape is determined by the form of the underlying bone. In common usage, the word nail often refers to this part only. The free margin (margo liber) or distal edge is the anterior margin of the nail plate corresponding to the abrasive or cutting edge of the nail [6]. The hyponychium (informally known as the "quick") is the epithelium located beneath the nail plate at the junction between the free edge and the skin of the fingertip. It forms a seal that protects the nail bed. The onychodermal band is the seal between the nail plate and the hyponychium. It is found just under the free edge, in that portion of the nail where the nail bed ends and can be recognized by its glassy, greyish colour (in fair-skinned people). It is not perceptible in some individuals while it is highly prominent on others. The eponychium is the small band of epithelium that extends from the posterior nail wall onto the base of the nail. Often and erroneously called the "proximal fold" or "cuticle", the eponychium is the end of the proximal fold that folds back upon itself to shed an epidermal layer of skin onto the newly formed nail plate. This layer of non-living, almost invisible skin is the cuticle

that "rides out" on the surface of the nail plate. Together, the eponychium and the cuticle form a protective seal [7].

The cuticle on the nail plate is dead cells and is often removed during manicure, but the eponychium is living cells and should not be touched. The perionych is the projecting edge of the eponychium covering the proximal strip of the lunula [8]. The nail wall (vallum unguis) is the cutaneous fold overlapping the sides and proximal end of the nail. The lateral margin (margo lateralis) is lying beneath the nail wall on the sides of the nail and the nail groove or fold (sulcus matricis unguis) are the cutaneous slits into which the lateral margins are embedded. The paronychium is the border tissue around the nail and paronychia is an infection in this area.

### **NAIL DISEASES & DISORDERS**

**Paronychia** infections of the nail fold can be caused by bacteria, fungi and some viruses. The proximal and lateral nail folds act as a barrier, or seal, between the nail plate and the surrounding tissue. If a tear or a break occurs in this seal, the bacteria can easily enter [9]. This type of infection is characterized by pain, redness and swelling of the nail folds. People who have their hands in water for extended periods may develop this condition, and it is highly contagious.

Pseudomonas bacterial infection can occur between the natural nail plate and the nail bed, and/or between an artificial nail coating and the natural nail plate. Many people have been led to believe that the classic 'green' discoloration of this type of infection is some type of mold. In actuality, mold is not a human pathogen. The discoloration is simply a by-product of the infection and is caused primarily by iron compounds. Pseudomonas thrives in moist places; it feeds off the dead tissue and bacteria in the nail plate, while the moisture levels allow it to grow. The aftereffects of this infection will cause the nail plate to darken and soften underneath an artificial coating. The darker the discoloration, the deeper into the nail plate layers the bacteria has traveled. If the bacteria have entered between the nail plate and the nail bed, it will cause the same discolorations and may also cause the nail plate to lift from the nail bed [10].

A fungal or yeast infection which results in Onychomycosis, can invade through a tear in the proximal and lateral nail folds as well as the eponychium. This type of infection is characterized by onycholysis (nail plate separation) with evident debris under the nail plate. It normally appears white or yellowish in color, and may also change the texture and shape of the nail. The fungus digests the keratin protein of which the nail plate is comprised. As the infection progresses, organic debris accumulates under the nail plate often discoloring it. Other infectious organisms may be involved, and if left untreated, the nail plate may separate from the nail bed and crumble off [11].

**Tinea Unguis**, or ringworm of the nails, is characterized by nail thickening, deformity, and eventually results in nail plate loss.

**Onychatrophia** is an atrophy or wasting away of the nail plate which causes it to lose its luster, become smaller and sometimes shed entirely. Injury or disease may account for this irregularity.

The aforementioned nail irregularities are among those I have been witness to during my years in the salon. There are others that only a trained dermatologist will be able to diagnose and treat. Some are contagious, and some are the result of injury or illness [12]. Physicians will sometimes examine your fingernails because many diseases will appear as various changes in the nail plate.

Any change in the nail plate could be cause for concern, whether it is a simple splinter hemorrhage that appears as a tiny black line in the nail plate, or a drastic change. Nail technicians are trained to beautify the hands/feet and are not allowed to diagnose nail diseases or to treat them in the salon. For your nail health, seek the diagnosis and recommendation of a knowledgeable dermatologist.

### **TREATMENTS FOR BRITTLE NAILS**

A lot of things can go wrong with nails. They can be brittle, pitted, discolored, flaky and ridged, among other common (and unpleasant) abnormalities. But brittle nails can be a natural part of the aging process, and they're often represented by dry, cracking or splitting nails. They may also grow slowly or simply seem weak and easily breakable. Nail abnormalities may be caused by a relatively minor condition like brittle nail syndrome, which is what it sounds like: excessively brittle nails often caused by a lack of moisture. They also could stem from a lack of iron or zinc. But nail problems may also be representative of something more severe, such as hepatitis, jaundice, lupus or heart disease. When in doubt, look for basic causes first, such as a fungal infection, a reaction to nail polish or bruising from an impact. There's ongoing debate about whether brittle nails are caused predominantly by a lack of protein or moisture in the nails. Consequently, most treatments for brittle nails are concerned with one of these two factors. In this article, we'll look at five ways of boosting nail health and learn why a candle may be just what your troublesome fingernails need [13].

### **Vitamin Supplements and Biotin;**

Vitamins are a key factor in making bodily processes run effectively and healthily, and nails are no exception. A lack of iron and zinc can harm nail health, and a basic multivitamin is often the solution. Try something with staples like niacin, iron, calcium and vitamins A and C, vitamin B complex containing biotin is often cited as important for nail health. Besides being present in certain vitamin supplements, biotin can be found

in oatmeal, bananas, mushrooms, peanuts, soy and, if you can stomach it, some animal organs. It's not really clear how effective biotin is in strengthening nails or by what mechanism it operates, though the vitamin has been successful in strengthening horse hooves. In one test, women who took 2.5 milligrams of biotin a day for six months or more ended up with 25 percent thicker nails. Biotin is found in many foods, so most people normally ingest enough, except in certain cases, such as people with alcoholism, people who eat excessive quantities of raw egg whites or those who use antibiotics for an extended period. Many pregnant women have a biotin deficiency, which can lead to birth defects, making prenatal vitamins essential [14]. Some healthy women who aren't pregnant take prenatal vitamins for their reputed benefits for hair and nail health. Much information about supplement-nail relationships is anecdotal rather than scientifically established. There is some evidence to suggest that glucosamine, often used for the treatment of osteoarthritis, is beneficial. Gelatin and an herb known as horsetail are often used for treating brittle nails, though there's little or conflicting evidence that they actually help (Gelatin is derived from animal hooves and connective tissue and is a favorite of many nail salons).

### **Super Moisturizers**

Often with brittle nails, the main culprit is simply a lack of moisture, just as dry skin can leave your epidermis cracked or flaky. Regular moisturizers available at the drugstore, such as Vaseline, can help to keep nails healthy, while some people trust home remedies, like a mix of egg yolks and milk. There are also creams that seal in moisturizers, such as Aquaphor and Trind Nail Balsam. Over the last decade, a class of creams called super moisturizers has become firmly entrenched in the nail care market. Applied to nails and the area surrounding them, super moisturizers are creams beefed up with vitamin E, avocado oil and shea butter [15-17].

### **Fortified Nail Polishes**

Nail polishes don't have to be simply cosmetic enhancers. Fortified nail polishes are packed with extra vitamins and minerals and promise to boost nail health. Some of them have rather ambitious names e.g., Sally Hansen Miracle Cure and equally lofty claims. (Consider again the Sally Hansen product, which cites laboratory data claiming 50 percent stronger nails in three days. But it's not just about what type of polish you put on. You should also pay attention to what you use to take off nail polish. Nail products, particularly nail polish removers, can contain some harmful ingredients. Avoid any products containing formaldehyde, acetone or toluene, all of which can harm nail health. Formaldehyde, the same ingredient used in embalming, and acetone can dry out nails. Camphor and phthalates may also cause allergic reactions.

### **The Natural Approach**

Avoid nail products that could cause an allergic reaction, like those with ingredients listed on the previous page. And don't trust nail hardeners: Brittle nails actually are already too hard, and nail hardeners produce allergic reactions in some people. You're better off moisturizing your nails. Keep your cuticles: They're actually important for nail health, and trimming them excessively can leave you more prone to an infection. Cutting them may also lead to nail deformities. If a hangnail or excessive cuticle is bothering you, use scissors to cut cleanly and moderately. Lay off the nail polish occasionally. It gives your nails a break, letting them breathe and allowing you to look at the physical appearance of your nails and make sure there aren't any issues lurking underneath the polish. Moisturizing creams will be better able to do their work on an unvarnished nail. You should also minimize the amount of nail polish remover you use, applying only as much as is necessary to remove the polish. As we said before, avoid removers containing harmful ingredients like formaldehyde, which dries out nails. Finally, it's good to let your nails get some air, but keep an eye on how they're affected by the environment. Cold, dry air can lead to cracking [18].

### **Protect Your Hands**

Protect your nails by protecting your hands. Wear gloves, particularly in cold weather or when washing dishes. Excessive hand washing allows water to seep into nails, swelling them and leading to brittleness. Keep nails short. They're more likely to be damaged when they become long, and they could get in the way of day-to-day tasks. Office work, in particular, can be hard on your hands, as it involves a lot of manual activity. Stay alert, such as by carefully closing drawers so as not to catch your fingers in them. Poor circulation may also be a contributing factor to brittle nails, so if you think you suffer from this condition, talk with your doctor. Other health problems, such as an underactive thyroid, may harm nail growth and should be discussed with a physician. Keep an eye out for signs of fungal infections, which are particularly common in senior citizens. Although these infections can be treated, because of ineffective creams or potential side effects from oral medications, some doctors recommend leaving a fungal infection alone and monitoring it [15].

### **FACTORS AFFECTING DRUG TRANSPORT INTO/ACROSS THE NAIL**

Topical application of a drug formulation on the nail plate, the drug has to enter the nail plate and diffuse into the deeper nail layers and possibly into the nail bed. Walters et al. found that the nail plate behaves like a concentrated hydro gel rather than a lipophilic membrane. Drug delivery into and through the nail plate is influenced by:

- Physicochemical properties of a drug molecule to be applied,
- Type and nature of formulations

- Presence of permeability enhancers in the formulations
- Properties nail and
- Interactions between the permeant and the keratin network of the nail plate.

#### **Molecular size of drug**

The larger the molecular size, the harder it is for drug to diffuse through the keratin network and lower the drug permeation. Mertin and Lippold demonstrate the decreasing permeability coefficients through human nail plate and through bovine hoof membrane with increasing molecular size of a series of alkyl nicotines.

#### **Hydrophilicity / lipophilicity of drug**

Walters et al. studied the permeation of a series of homologous alcohols (C1–C12), diluted in saline, through avulsed human nail plates. Increasing the chain length from one carbon to eight carbon atoms resulted in a decrease in permeability coefficient, after which, increasing chain length (>C12) resulted in increased permeability coefficient. The study concludes that the nail plate is characterized as a hydrophilic gel membrane [12].

#### **Nature of Vehicle used in formulation**

The permeability coefficients of alcohols diluted in saline through nail plates was five times greater than the permeability coefficients of neat alcohols. Water hydrates the nail plate which consequently swells. Considering the nail plate to be a hydrogel, swelling results in increased distance between the keratin fibres, larger pores through which permeating molecules can diffuse and hence, increased permeation of the molecules. Replacing water with a non-polar solvent, which does not hydrate the nail, is therefore expected to reduce drug permeation into the nail plate [18].

#### **pH of vehicle and solute charge**

The pH of aqueous formulations affects the ionization of weakly acidic/basic drugs, which in turn influences the drug's hydrophilicity /hydrophobicity, solubility in the drug, formulation, solubility in the nail plate and its interactions with the keratin matrix. It seems that the pH of the formulation has a distinct effect on drug permeation through the nail plate.

#### **ELECTROCHEMOTHERAPY FOR NAIL DISORDERS**

The goal of this therapy is to develop an active method of drug delivery across the nail plate which in turn is believed to increase the success rate of topical monotherapy and decrease the duration of treatment of nail disorders. Currently the electrically mediated techniques for drug delivery across the nail plate are investigated. Recently the iontophoresis nail delivery method studied. Iontophoresis was found to enhance the transport of drugs

across the nail plate significantly. Similar to transdermal iontophoresis, the predominant mechanisms contributing to enhanced transport of drugs in the case of trans nail iontophoresis are electrophoresis and electro osmosis [15].

#### **MESOSCISSIONING TECHNOLOGY**

MesosciSSIONING technology creates a Micro-conduit through the skin or nail within a specified depth range. Fully open pathways can be painlessly scized (cut) through the stratum corneum of the skin or through the nail. Micro conduits, 300-500 microns in diameter, are produced within seconds and without sensation. These pathways can be used to deliver drugs across the skin. Such micro conduits also permit access for sub dermal analyte extraction (including blood for glucose testing). In addition, they reduce the skin electrical impedance to less than 1000 ohms for biopotential measurements. In nails, micro conduits quickly reduce the painful pressure of sub ungula hematoma (black toe) and could serve as a prophylactic to prevent such pressure build-up in runner's nails.

#### **NANOPATCH NAIL FUNGUS**

Nano Patch Fungus uses AC/DC electro chemistry and targeted drug delivery to actively push Anti-fungal drugs right through the nail cuticle to the actual location of the fungus growth. This would be the first treatment option to directly target nail fungus at its source of growth.

#### **IONTOPHORETIC DRUG DELIVERY ACROSS HUMAN NAIL**

Topical trans-nail delivery of antifungal drugs is limited by several physicochemical and physiological factors. Use of chemical permeation enhancers has been a common approach for enhancing trans-nail delivery of drugs. The potential of physical permeation enhancement techniques has been found to be higher than the potential of chemical permeation enhancers in transdermal delivery of hydrophilic drugs and macromolecular therapeutic agents. However, application of physical permeation enhancement techniques has not been explored for trans-nail drug delivery. In the current work, iontophoresis was applied across human nail *in vitro* to assess its efficiency in enhancing drug delivery. Salicylic acid (SA) was used as test diffusant. The influence of pH, ionic strength, and current density was studied. Obviously, increase in current density increased the trans-nail transport flux. It appears that about 50-100mM ionic strength is required for optimal conduction of electric current across nail. The flux enhancement factor (iontophoretic flux/passive flux) also increased with increase in pH due to increased ionization of SA. This study demonstrates the efficacy of iontophoresis in enhancing the trans-nail delivery of drugs. Topical drug delivery to treat nail diseases such as onychomycosis and psoriasis is receiving increasing attention. Topical nail delivery is challenged by the


complicated structure of the nail and the low permeability of most drugs across the nail plate. Considerable effort has been directed at developing methods to promote drug permeation across the nail plate. Iontophoresis efficiently enhances molecular transport across the skin and the eye and is now being tested for its potential in unguinal delivery [18].

**NOVEL DRUG-LOADED PATCHES FOR THE TREATMENT OF NAIL DISEASES**

Nail diseases are common and require treatment. For example, nail fungal infections, affect up to 40% of the population, are more common in the elderly, immune suppressed and diabetics and can significantly affect the quality of life of sufferers. For example, walking is painful due to unsightly nails inhibit social interactions and work. The main treatment regimen for fungal infections - oral therapy with anti fungals suffers from serious drawbacks such as liver toxicity. Another common nail disease, psoriasis, is treated with repeated and extremely painful injections of the drug into the skin surrounding the nail. Effective topical therapy could replace or complement the current and thereby lead to elimination/ of the adverse effects of oral/ injected therapies. Unfortunately, the existing topical products for fungal infections have shown limited success, and no topical products have been licensed for nail psoriasis. Research into the topical therapy of nail diseases is therefore essential to address this unmet clinical need. We propose to investigate patches as drug carriers for the topical treatment of nail diseases. Following application to the nail plate, the patch would remain at the disease site and continuously release drug for long durations [19]. This non-invasive method of delivery is expected to be popular with patients. Patches are already commercially available for application to the skin.

However due to the differences between the drugs, and between the skin and the nail surfaces, skin patches cannot simply be loaded with drugs for nail diseases. Nail patches have to be formulated from scratch. Our aim is to develop a method for the rational design of drug-loaded nail patches. By investigating the underpinning science, and by exploiting our combined expertise in engineering, modelling, and drug delivery, and the facilities at our two institutions, we will develop a scientific method to formulate nail patches which will be universally applicable to a range of diseases and drugs. Thus, the proposed work is fundamentally different from those described in the scarce literature on nail patches, where the latter have only been used as a convenient vehicle, or where the influence of important patch components on patch efficacy was tested in an empirical manner. We will use a combination of theoretical, experimental and modelling methods to: i) identify the correct patch components, such as the adhesive, backing membrane, solvents, ii) formulate drug-loaded nail patches, and iii) evaluate the formulated patches in terms of adhesion to the nail plate, effects on nail plate hydration, drug transfer in to the nail plate, and subsequent drug action against the disease. Theoretical concepts such as solubility parameter, and finite element analysis and cohesive zone models will be used to select the correct parameters, reduce the number of required experiments, and ultimately provide some ready-to-use tools. Assessment of the efficacy of the formulated patch against a nail disorder will indicate the success of the developed methodology. To enable the testing of patch efficacy, fungal infection is chosen as the nail disorder, and the anti-fungal drugs, terbinafine and amorol fine will be used in this work. Use of two drugs will give an indication of the robustness and universality of the developed methods [20].


**Figure 1. Different parts of nail**




**Figure 3. Paronychia Infection**



**Figure 2. Semi Diagrammatic Cross Section of Finger Nail**



**Figure 4. Pseudomonas bacterium trapped between the nail plate and the nail bed. (The 'Greenies')**



**Figure 5. Fungal Infection of the nail plate**



**Figure 6. Ringworm of the nails**



**Figure 7. Ingrown Toenail**



## DISCUSSION

Topical drug delivery is especially suitable for onychomycosis (fungal infections of the nail plate and/ or nail bed) and nail psoriasis, which affect 2- 13 and 1 - 3% of the general population, respectively, and make up the bulk of nail disorders. Topical therapy would avoid the adverse events and drug interactions of systemic antifungal agents and the pain of injection when anti-psoriatic agents are injected into affected nail folds. However, successful topical therapy is extremely challenging due to the very low permeability of the nail plate. Five speakers spoke about various aspects of topical drug delivery to the nail, including review of the nail plate structure, function, diseases, their existing therapies (systemic and topical), limitations and global sales. The need for effective topical drug delivery to the nail to overcome the problems associated with present treatment, and the fact that there are few topical formulations available for the treatment of nail fungal infections and psoriasis, and the even fewer effective formulations, was highlighted. The absorption of drugs into the nail unit, following topical application to the nail plate, is highly desirable to treat nail disorders, such as onychomycosis (fungal infections of the nail). Nail permeability is however quite low and limits topical therapy to early/mild disease states. In this paper, the recent research into unguinal drug delivery is reviewed. The nail unit and the two most common diseases affecting the nail-onychomycosis and nail psoriasis--are briefly described to set the scene and to give an overview of the nature and scope of the problem. The factors, which affect drug uptake and permeation through the nail plate such as solute molecular size, hydrophilicity/ hydrophobicity, charge, and the nature of the vehicle, are then discussed, followed by ways of enhancing drug transport into and through the nail plate. Finally, drug-containing nail lacquers which, like cosmetic varnish, are brushed onto the nail plates to form a film, and from which drug is released and penetrates into the nail. The permeability of topically applied drugs through keratinized nail plate is highly poor

## REFERENCES

1. Vajnovic I, Simmler, L, Betz G. Investigation of different formulations for drug delivery through the nail plate. *International Journal of Pharmaceutics*, 386, 2010, 185–194.
2. Gunt HB and Kasting GB. Effect of hydration on the permeation of ketoconazole through human nail plate in vitro. *European Journal of sciences*, 32, 2007, 254–260.

and drug uptake into the nail apparatus is extremely low. Nail lacquers containing drugs are an innovative type of dosage form. Like cosmetic nail varnish, they are applied on to the nail plate using a brush. The field of unguinal drug delivery following topical application is not fully explored and more research in this field is needed to resolve the conflicting reports on the physicochemical parameters that influence unguinal drug permeation and to find and characterize new penetration enhancers and delivery vehicles. Drug transport into nail plate is assisted by filing the nail plate before topical application of drug formulations as well as by the use of chemical enhancers. The oral therapies encounter side effects and topical therapies for nail diseases are limited by poor permeability of nail plate. An optimal penetration enhancer would improve drug delivery through nail plate facilitating new possibilities for treating neighboring target sites if the systemic circulation is reached.

## CONCLUSION

Drug delivery to the nail (unguinal drug delivery) constitutes a major challenge, with the lack of understanding of both the barrier properties of the nail and formulations to achieve enhanced unguinal delivery restricting the efficiency of topical treatments for nail disorders. Topical delivery of systemic therapeutics offers benefits but presents a greater technical challenge. Among the benefits, first pass avoidance, convenience and sustained release are most often sited. Nail diseases like onychomycosis, nail psoriasis, yellow nail syndrome, paronychia and many more, being cured successfully using medicated lacquers. This avoids the oral toxicity of anti-fungal drugs and provides longer contact time at the site of action. This systemic review covers the anatomy of a human nail, diseases related to nail plate, the formulations designed for nail application and some techniques used to enhance the topical bioavailability of the drugs across the nail, latest trends in drug delivery across the nail.

3. Rubio MC, Ariz IR, Gil J, Benito J, Rezusta A. Potential fungicidal effect of Voriconazole against *Candida* spp. *International Journal of Antimicrobial Agents*, 25, 2005, 264–267.
4. Berker DA, Andre J, Baran R. Nail biology and nail science. *International Journal of Cosmetic Science*. 2007; 29:241-275.
5. Lewin K. The normal finger nail. *British J.Dermatology*, 77, 1961, 485-87.
6. Dykyj JD. *Anatomy of the nail clin padiatr.med.surgery*, 6(2), 1989, 21-28.
7. Forslind B. On the structure of the normal nail.A scanning electrion microscopy study. *Archives for dermatologishe for schung*, 251, 1975, 199.
8. Kobayashi Y, Miyamoto M, Sugibayashi K, Morimoto Y. Drug permeation through the three layer of the of the human nail plate. *J.Pharma Pharmacol*, 51, 1999, 271-78.
9. Hashimoto K, Bernard GG, Nelson R, Lever WF. *J Invest. Derm*, 47, 1966, 205-217.
10. Kenneth A, Walters, Gordon L, Flynn, &John R.Marvel. Physiochemical characterization of the human nail. *J. Pharm. Pharmacol*, 35, 1983 , 28-33
11. Runne U. The human nail; structure, growth &pathological changes. *Current problem in dermatology*, 9, 1981, 102.
12. Baran R, Dawber RPR, eds. Diseases of the nail& their management, Oxford: Blackwell scientific publication, 1984.
13. Sau P et al. Brwens disease of the nail bed & periungnal area. *Arch dermatol*, 130, 1994, 204.
14. Gauri Malhotra, Joel L. Zatz Investigation of nail permeation enhancement by chemical modification. *Journal of Pharm .Sci*, 91(2), 2002, 312-323.
15. Sun Y, Liu JC, Kimbleton ES, Wang JCT. Antifungal treatment of nails, United states patent US 0056164A, 1997.
16. Soong MH. Transport properties of drugs& model compounds across the human nail. PhD dissertation. University of Minnesota, 1991.
17. Robbins CR. Chemical & Physical behaviour of human nail,3rd edition, New York: Springerverlag, 1997, pp.93-130
18. Heel RC, Brogden RN, Speight TM. Econazole; A review of its antifungal activity and therapeutic efficacy. *Drug*, 16, 1978, 177-201.
19. Gupchup CV, Zatz JL. Structural characteristics and permeability properties of human nail- a review. *J.Cosmet.Sci*, 1999, 363-385.
20. JU-H Kim, Chi H Lee. A method to measure the amount of drug penetrated across the nail plate. *Pharma Research*, 18(10), 2001, 1468-1471.